

Final approval of the article: RD, JW, AP, AZ, BS, DW

Statistical analysis: RD, JW

Obtained funding: Not applicable

Overall responsibility: RD

REFERENCES

1. Kearon C. Natural history of venous thromboembolism. *Circulation* 2003;107(23 Suppl 1):I22-30.
2. Righini M, Bounameaux H. Clinical relevance of distal deep vein thrombosis. *Curr Opin Pulm Med* 2008;14:408-13.
3. Lagerstedt CI, Olsson CG, Fagher BO, Oqvist BW, Albrechtsson U. Need for long-term anticoagulant treatment in symptomatic calf-vein thrombosis. *Lancet* 1985;2:515-8.
4. Pellegrini VD Jr, Langhans MJ, Totterman S, Marder VJ, Francis CW. Embolic complications of calf thrombosis following total hip arthroplasty. *J Arthroplasty* 1993;8:449-57.
5. Labropoulos N, Kang SS, Mansour MA, Giannoukas AD, Moutzourous V, Baker WH. Early thrombus remodelling of isolated calf deep vein thrombosis. *Eur J Vasc Endovasc Surg* 2002;23:344-8.
6. Wang CJ, Wang JW, Weng LH, Hsu CC, Lo CF. Outcome of calf deep-vein thrombosis after total knee arthroplasty. *J Bone Joint Surg Br* 2003;85:841-4.
7. Oishi CS, Grady-Benson JC, Otis SM, Colwell CW Jr, Walker RH. The clinical course of distal deep venous thrombosis after total hip and total knee arthroplasty, as determined with duplex ultrasonography. *J Bone Joint Surg Am* 1994;76:1658-63.
8. Solis G, Saxby T. Incidence of DVT following surgery of the foot and ankle. *Foot Ankle Int* 2002;23:411-4.
9. Higgins J, Green S. *Cochrane handbook for systematic reviews of interventions*, version 5.0.2 [updated 2009]. The Cochrane Collaboration; 2009.
10. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med* 2009;151:W65-94.
11. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008-12.
12. Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses.
13. Nielsen HK, Husted SE, Krusell LR, Fasting H, Charles P, Hansen HH. Silent pulmonary embolism in patients with deep venous thrombosis. Incidence and fate in a randomized, controlled trial of anticoagulation versus no anticoagulation. *J Intern Med* 1994;235:457-61.
14. Dorr LD, Gendelman V, Maheshwari AV, Boutary M, Wan Z, Long WT. Multimodal thromboprophylaxis for total hip and knee arthroplasty based on risk assessment. *J Bone Joint Surg Am* 2007;89:2648-57.
15. Masuda EM, Kessler DM, Kistner RL, Eklof B, Sato DT. The natural history of calf vein thrombosis: lysis of thrombi and development of reflux. *J Vasc Surg* 1998;28:67-73; discussion 474.
16. Solis MM, Ranval TJ, Nix ML, Eidt JF, Nelson CL, Ferris EJ, et al. Is anticoagulation indicated for asymptomatic postoperative calf vein thrombosis? *J Vasc Surg* 1992;16:414-8; discussion 418-9.
17. Lohr JM, James KV, Deshmukh RM, Hasselfeld KA, Allastair B. Karmody Award. Calf vein thrombi are not a benign finding. *Am J Surg* 1995;170:86-90.
18. Sachdev U, Teodorescu VJ, Shao M, Russo T, Jacobs TS, Silverberg D, et al. Incidence and distribution of lower extremity deep vein thrombosis in rehabilitation patients: implications for screening. *Vasc Endovasc Surg* 2006;40:205-11.
19. Pinede L, Ninet J, Duhaut P, Chabaud S, Demolombe-Rague S, Durieu I, et al. Comparison of 3 and 6 months of oral anticoagulant therapy after a first episode of proximal deep vein thrombosis or pulmonary embolism and comparison of 6 and 12 weeks of therapy after isolated calf deep vein thrombosis. *Circulation* 2001;103:2453-60.
20. Philbrick JT, Becker DM. Calf deep venous thrombosis. A wolf in sheep's clothing? *Arch Intern Med* 1988;148:2131-8.
21. Righini M. Is it worth diagnosing and treating distal deep vein thrombosis? No. *J Thromb Haemost* 2007;5(Suppl 1):55-9.
22. Righini M, Paris S, Le Gal G, Laroche JP, Perrier A, Bounameaux H. Clinical relevance of distal deep vein thrombosis. Review of literature data. *J Thromb Haemost* 2006;9:56-64.
23. Hogg K, Ashton A. Towards evidence based emergency medicine: best BETs from the Manchester Royal Infirmary. Antithrombotic treatment of below knee deep venous thrombosis. *Emerg Med J* 2003;20:364-5.
24. Geerts WH, Bergqvist D, Pineo GF, Heit JA, Samama CM, Lassen MR, et al. Prevention of venous thromboembolism: American College of Chest Physicians evidence-based clinical practice guidelines (8th edition). *Chest* 2008;133(6 Suppl):381-453S.
25. Schulman S, Beyth RJ, Kearon C, Levine MN. Hemorrhagic complications of anticoagulant and thrombolytic treatment: American College of Chest Physicians evidence-based clinical practice guidelines (8th edition); *Chest* 2008;133(6 Suppl):257-98S.
26. Saarinen JP, Domonyi K, Zeitlin R, Salenius JP. Postthrombotic syndrome after isolated calf deep venous thrombosis: the role of popliteal reflux. *J Vasc Surg* 2002;36:959-64.
27. McLafferty RB, Moneta GL, Passman MA, Brant BM, Taylor LM Jr, Porter JM. Late clinical and hemodynamic sequelae of isolated calf vein thrombosis. *J Vasc Surg* 1998;27:50-6; discussion 56-7.
28. Meissner MH, Caps MT, Bergelin RO, Manzo RA, Strandness DE Jr. Early outcome after isolated calf vein thrombosis. *J Vasc Surg* 1997;26:749-56.

Submitted Jun 27, 2011; accepted Sep 24, 2011.

Additional material for this article may be found online at www.jvascsurg.org.

DISCUSSION

Dr Joseph Ricotta (*Atlanta, Ga*). How long were these patients anticoagulated for? Was there variability between the studies in terms of the duration of anticoagulation therapy?

Dr Randall R. De Martino. There was some variation. We required a minimum of 30 days because most of these were postoperative patients and that is what reporting goes to, although I think treatment should last for at least 6 weeks. We did a sensitivity analysis of outcomes by studies reporting 6 weeks of anticoagulation treatment or less. There was no change in the effect on clot propagation or PE.

Dr Ricotta. The longer the duration?

Dr De Martino. Correct.

Dr James Debord (*Peoria, Ill*). Do you think your analysis and results can be extrapolated to isolated gastrocnemius and

soleus plexus clots in the absence of the named tibial vessel clots?

Dr De Martino. We did not look specifically at gastrocnemius or soleal clots in this study. We were focused on tibial and peroneal thrombi. I think that they may follow a similar pattern, although we can't make any conclusions based on this study. Other studies that have addressed that question and maybe a compilation of those studies can help identify if they are similar in characteristics. But I don't know that the clinical history of them is as well defined as is tibial and peroneal.

Dr Firas Mussa (*New York, NY*). Your last bullet suggests future studies. Can you elaborate on design of those studies?

Dr De Martino. I think there is still clinical equipoise as to whether or not we can treat or not treat these patients. Based on

the data and the studies that are available, I think that there is still a debate about this. I think clinical randomized trials will need to assess the benefits and the harms of treatment, including bleeding, mortality rates, and post-thrombotic syndrome. I am aware of one ongoing clinical study in Europe, but I am unaware of the current enrollment status. I was told yesterday in discussion, the University of Washington has applied for a grant to perform a randomized trial for this, and we hope that that will be funded and shed light on this topic.

Dr Cynthia Shortell (*Durham, NC*). It is no surprise that anticoagulation prevents thrombotic complications of calf vein clot. The question has always been, when is the risk of anticoagulation worth the benefit? And in calf vein thrombosis, that issue gets down to what the risks of those patients are. One would intuit that high-risk patients with calf clots, such as neurosurgical patients or immobile patients, would be at higher risk, but that's unknown at this point. Did any of the studies that you included contain that information? And if so, do you think you could use it to help us identify those patients with calf clot that would really benefit from anticoagulation?

Dr De Martino. I think we agree that there are patients who are at higher risk. Several of these studies did try to identify specific risk factors that would identify those patients with thrombi that are going to propagate or cause PE. The most consistent risk factor identified was inpatient status. Many other patient-level variables that they were able to analyze were not predictive in their analyses. We didn't include that type of review in the present study, but that is what I have been able to extract from the data that is available to date.

Dr Mark Meissner (*Seattle, Wash*). I was somewhat surprised that although bleeding was heterogeneous, pulmonary embolism and thrombus propagation was remarkably homogeneous. I think this is surprising, because perhaps even more so than with proximal vein thrombosis, calf vein thrombosis is often perceived to be a heterogeneous disease with the risk of complications based on the underlying risk factors for thrombosis. Your analysis seems to imply that all calf vein thrombi should be treated as equal, which may not be correct. Can you explain the homogeneity of your findings?

Dr De Martino: In regards to PE and clot propagation, I think our pooled summary was homogeneous because of our strict entry criteria. There are other studies reported on calf DVT that were excluded because they didn't meet this criteria. I think that helped in creating a homogeneous estimate.

The majority of studies did report on PE and clot propagation. Very few studies reported bleeding and mortality complications, so the data that we were able to extract were sparse and spread across eight studies, where only two to three may have contributed to the data, and I think that that affected our ability to make a homogeneous summary effect.

I think that while we were able to make a homogeneous study effect, we acknowledge that the strength of our findings comes from the quality of the studies that they were derived from. And so a robust clinical trial will really answer this question in the best light possible.

Dr Peter Gloviczki (*Rochester, Minn*). Did you notice a difference in symptomatic and asymptomatic calf vein thrombosis, or were all these patients symptomatic?

Dr De Martino. We didn't perform a formal subgroup analysis of studies that were symptomatic or asymptomatic. Approximately half of the studies identified their patients as symptomatic, two of those four being the randomized controlled trials. But we are unable to make a comment about whether the symptomatic status does make a difference.

Dr Gloviczki. And was the diagnosis of PE made using similar techniques in these studies?

Dr De Martino. The diagnosis of PE was similar in all but one study, which used a combination of V/Q scan and clinical diagnosis based on patients with a sudden death that was preceded by rapid leg swelling. And so it was felt that that was a clinical diagnosis of PE and we did include that.

Dr Roger Shinnerl (*Evansville, Ind*). I was wondering if you could please tell me if I am wrong, but I was left with the distinct impression, after looking at your data briefly, that all of the positive results were out of one study, the Pellegrini study. And that disturbs me. Does that not make that study an outlier? And even if not, should we really be endorsing that finding with the results of a pooled analysis?

Dr De Martino. That one study was a nested case-control study within a randomized trial. While they did identify a lot of outcomes, I do not think that their methods were any different than the other randomized trial that would make me think that they should be excluded or be an outlier. I think the fact that our summary of facts remained homogeneous despite their inclusion attests that it was appropriate to include them within our analysis.